

SARS-CoV-2 UV Dose-Response Behavior

White Paper Prepared for IUVA by:

Ernest R. Blatchley III,^a Brian Petri,^b Wenjun Sun^c

^aLee A. Rieth Professor in Environmental Engineering, Purdue University (blatch@purdue.edu)

^bResearch Director, Trojan Technologies (bpetri@trojanuv.com)

^cAssociate Professor in School of Environment, Tsinghua University (wsun@tsinghua.edu.cn)

Ultraviolet (UV) radiation is a broad-spectrum antimicrobial agent that has been applied successfully in a wide range of disinfection applications. UV radiation in the wavelength range $200\text{ nm} \lesssim \lambda \lesssim 320\text{ nm}$, sometimes referred to as “germicidal” or “microbicidal” UV radiation, is known to cause damage to DNA and RNA that results in inactivation of microorganisms and viruses. For radiation with wavelengths less than about 240 nm, damage to proteins can also contribute to inactivation. Given that all viruses contain a nucleic acid molecule, either DNA or RNA, and a protein coat (called a capsid) that surrounds the nucleic acid, all viruses are susceptible to inactivation by exposure to UV. However, viral sensitivity to UV is quite variable, though the development of a comprehensive understanding of the causative factors is still an active area of research.

SARS-CoV-2 (aka the novel coronavirus) is the virus that causes Coronavirus Disease 2019 (COVID-19). Transmission of COVID-19 appears to be largely associated with airborne particles that may be released by symptomatic or asymptomatic individuals who have been infected by SARS-CoV-2,¹ although it is also known that the virus can remain infective on surfaces for as much as 24-72 hours, depending on the material it is contact with,² so contact with surfaces represents another possible mechanism of disease transfer.³

At present, only limited data are available to define how SARS-CoV-2 responds to common disinfectants, including UV radiation. The standard method for defining the response of a microorganism to UV radiation is to measure its UV dose-response behavior, which describes the kinetics of inactivation. UV dose-response behavior represents a key piece of information for the design of UV disinfection systems.

Previous research has allowed definition of UV dose-response behavior for other, closely related viruses. To date, three studies have been published that have reported UV dose-response behavior of SARS-CoV (see Figure 1); this is the virus that caused an epidemic of Severe Acute Respiratory Syndrome (SARS) that affected roughly 8000 people in 26 countries in 2003.⁴ All of these studies were based on UV radiation at or near the wavelength (254 nm) that characterizes the output of low-pressure (LP) mercury lamps, which are the most commonly-used source of germicidal UV radiation. SARS-CoV and SARS-CoV-2 are structurally-similar viruses; both are non-segmented, enveloped, single-stranded RNA (ssRNA) viruses. It is likely that the UV dose-response behaviors of SARS-CoV and SARS-CoV-2 will be similar, though it is important to note that this has not yet been proven. Note also that there is considerable variability in the reported UV dose-response behavior of SARS-CoV. The variability that is evident in Figure 1 is probably due to deficiencies in the experimental methods that were used in some or all of these previous studies. The stated fluences (doses) are probably overestimates as the UV absorbance of the suspensions were not reported or explicitly accounted for; similarly, it was not clear that the conditions of UV exposure would allow for accurate calculation of the UV dose applied to the viral suspension. Based on the information provided in the reported studies of SARS-CoV UV dose-response behavior, the data set of Kariwa *et al.* appears to represent the most accurate, though it may also show false-high resistance.

Viruses are often characterized according to the presence/absence of an envelope, the form of their nucleic acid (single-stranded [ss] or double-stranded [ds]), and the type of nucleic acid (RNA or DNA);

SARS-CoV-2 is an enveloped, ssRNA virus. The form of the genome may play an important role in governing the sensitivity of viruses to UV exposure.^{5,6} Emerging models based on nucleotide sequences have a strong basis with known UV disinfection mechanisms, and may yield accurate predictions of the UV sensitivity of viruses.

Also included in Figure 1 are examples of the measured UV_{254} dose-response behavior of coliphage MS2, a non-enveloped ssRNA virus, and $\Phi 6$, an enveloped double-stranded RNA (dsRNA) virus. These two viruses have similar UV sensitivity as the SARS-CoV of Kariwa *et al.* (2004). These data suggest that if the UV sensitivity of SARS-CoV-2 is similar to the Kariwa *et al.* (2004) data, MS2 and $\Phi 6$ may represent suitable non-pathogenic surrogates.




Figure 1. Reported UV_{254} dose-response data for SARS-CoV in aqueous suspension.⁷⁻⁹ Note that the single data point provided by Duan *et al.* (2003) was based on radiation at 260 nm. Also included is a representative UV_{254} dose-response data set for coliphage MS2 in aqueous suspension (data courtesy of HDR/HydroQual) and a dose-response curve for Pseudomonas virus $\Phi 6$ from Ye *et al.* (2018).¹⁰

The effectiveness of UVC radiation as a disinfectant is influenced by the wavelength of radiation. This behavior has become increasingly important in recent years with the development of alternatives to conventional LP mercury lamps, including UV LEDs and plasma (excimer) lamps. The wavelengths of radiation produced by these sources depend on the chemical composition of the LED or the excimer. Collectively, these new sources provide access to radiation from across the germicidal UV spectrum.

A common graphical method for describing the effects of wavelength on microbial inactivation is the so-called “action spectrum.” In most cases, the action spectrum illustrates the relative rate of inactivation of

a microbe or virus at a given wavelength compared to its inactivation rate in response to irradiation at 254 nm. An example of a normalized action spectrum for MS2 in aqueous suspension is presented in Figure 2. As with most action spectra, the information presented in Figure 2 indicates that for wavelengths in the range 240–300 nm, peak inactivation efficiency is observed at about 265 nm, with steady decreases at wavelengths above and below this peak. In this wavelength range, the majority of viral inactivation is attributable to photochemical damage to its nucleic acid. For radiation at wavelengths less than about 240 nm, a rapid increase in the efficiency of inactivation is observed; this is attributable to damage to proteins, which is known to take place at these short wavelengths. It is expected that SARS-CoV-2 will display similar trends, but insufficient data are available at present to confirm or refute this hypothesis. The ultimate disinfection efficacy observed in an actual application will be influenced by the absorbance characteristics of the medium that is being disinfected. In some settings, there could be substances that could absorb strongly at wavelengths below 240 nm, which will mitigate the contributions of short wavelength UVC radiation.




Figure 2. Normalized action spectrum for coliphage MS2 (figure from Beck et al., 2015).¹¹ For all three data sets presented in this graph, the data were normalized against the measured response at 254 nm. Also included in this figure are action spectra presented from Rauth (1965) and Mamane-Gravetz (2005).^{12, 13}

The majority of UV dose-response data have been reported for experiments wherein the target virus was suspended in water. These experiments are critical for UV disinfection of water; however, the mechanism of transfer of SARS-CoV-2 (and many other viruses) generally involves viruses that are suspended in air or attached to surfaces. For both conditions, the virus itself is not suspended in water and will experience drying (desiccation). Desiccation, which will result from exposure to air, and will be influenced by relative humidity (RH), is known to represent a form of stress for most microbes and viruses, and may alter their sensitivity to other forms of environmental stress, including UV exposure, as illustrated in Figure 3.¹⁴⁻¹⁸




Figure 3. UV_{254} dose-response behavior of coliphage MS2 on surfaces (at two different values of RH) and in aqueous suspension. Data for MS2 in surfaces from Tseng and Li (2007).¹⁸ Data for MS2 in aqueous suspension were provided by HDR/HydroQual.

During the period of development of this white paper, several early reports of responses of SARS-CoV-2 and related viruses to UV radiation were reported in the scientific literature and in press releases. A study conducted by scientists at the U.S. Department of Homeland Security indicated that UVB radiation in ambient sunlight at irradiance values that are representative of mid-latitude, mid-day summertime conditions will result in 1 \log_{10} unit of inactivation in a time period of 7-15 minutes.¹⁹ A preprint publication by Bianco *et al.* (2020) reported that a UV_{254} dose of 3.7 mJ/cm^2 accomplished 3 \log_{10} units of SARS-CoV-2 inactivation when the virus was suspended in water (see <https://www.medrxiv.org/content/10.1101/2020.06.05.20123463v2.full.pdf>). A separate preprint document by Inagaki *et al.* indicated that a UV_{280} dose of roughly 38 mJ/cm^2 accomplished 3 \log_{10} units of inactivation of SARS-CoV-2 (see <https://www.biorxiv.org/content/10.1101/2020.06.06.138149v1>). Buonanno *et al.* (2020) reported that far-UVC (*i.e.*, wavelengths shorter than approximately 225 nm) accomplished at least 3 \log_{10} units of inactivation of airborne human coronaviruses alpha HCoV-229E and beta HCoV-OC43.²⁰

Collectively, these results are promising; however, it should be noted that some important details of these experiments were not reported in these documents, specifically relating to methods used to quantify dose delivery to samples containing the target virus. It is likely that publications in this area will continue to emerge, as well as our collective understanding of the effectiveness of germicidal UV radiation for control of SARS-CoV-2.

To summarize, available evidence suggests that UV radiation should be effective for inactivation of SARS-CoV-2 and other microbial pathogens. UV-based systems have important roles to play in battling SARS-CoV-2 in air, on surfaces, and in water. However, like all common disinfectants (*i.e.*, UV, chlorine, ozone), a need exists to quantify the kinetics of inactivation for SARS-CoV-2 for these applications.

References

1. Prather, K. A.; Wang, C. C.; Schooley, R. T., Reducing transmission of SARS-CoV-2. *Science* **2020**, *10.1126/science.abc6197*.
2. van Doremalen, N.; Bushmaker, T.; Morris, D. H.; Holbrook, M. G.; Gamble, A.; Williamson, B. N.; Tamin, A.; Harcourt, J. L.; Thornburg, N. J.; Gerber, S. I.; Lloyd-Smith, J. O.; de Wit, E.; Munster, V. J., Aerosol and Surface Stability of SARS-CoV-2 as Compared with SARS-CoV-1. *New England Journal of Medicine* **2020**, *382*, (16), 1564-1567.
3. Marshall III, W. F. Can COVID-19 (coronavirus) spread through food, water, surfaces and pets? <https://www.mayoclinic.org/diseases-conditions/coronavirus/expert-answers/can-coronavirus-spread-food-water/faq-20485479>
4. WHO SARS (Severe Acute Respiratory Syndrome). <https://www.who.int/ith/diseases/sars/en/>
5. Qiao, Z.; Ye, Y. Y.; Chang, P. H.; Thirunarayanan, D.; Wigginton, K. R., Nucleic Acid Photolysis by UV254 and the Impact of Virus Encapsulation. *Environmental Science & Technology* **2018**, *52*, (18), 10408-10415.
6. Kowalski, W. J.; Bahnfleth, W. P.; Hernandez, M. T. In *A Genomic Model for Predicting the Ultraviolet Susceptibility of Viruses and Bacteria*, IUVA World Congress, Amsterdam, Netherlands, 2009; Amsterdam, Netherlands, 2009.
7. Kariwa, H.; Fujii, N.; Takashima, I., Inactivation of SARS coronavirus by means of povidone-iodine, physical conditions, and chemical reagents. *Japanese Journal of Veterinary Research* **2004**, *52*, (3), 105-112.
8. Duan, S. M.; Zhao, X. S.; Wen, R. F.; Huang, J. J.; Pi, G. H.; Zhang, S. X.; Han, J.; Bi, S. L.; Ruan, L.; Dong, X. P.; Team, S. R., Stability of SARS coronavirus in human specimens and environment and its sensitivity to heating and UV irradiation. *Biomedical and Environmental Sciences* **2003**, *16*, (3), 246-255.
9. Darnell, M. E. R.; Subbarao, K.; Feinstone, S. M.; Taylor, D. R., Inactivation of the coronavirus that induces severe acute respiratory syndrome, SARS-CoV. *Journal of Virological Methods* **2004**, *121*, (1), 85-91.
10. Ye, Y. Y.; Chang, P. H.; Hartert, J.; Wigginton, K. R., Reactivity of Enveloped Virus Genome, Proteins, and Lipids with Free Chlorine and UV254. *Environmental Science & Technology* **2018**, *52*, (14), 7698-7708.
11. Beck, S. E.; Wright, H. B.; Hargy, T. M.; Larason, T. C.; Linden, K. G., Action spectra for validation of pathogen disinfection in medium-pressure ultraviolet (UV) systems. *Water Research* **2015**, *70*, 27-37.
12. Rauth, A. M., PHYSICAL STATE OF VIRAL NUCLEIC ACID AND SENSITIVITY OF VIRUSES TO ULTRAVIOLET LIGHT. *Biophysical Journal* **1965**, *5*, (3), 257-&.
13. Mamane-Gravetz, H.; Linden, K. G.; Cabaj, A.; Sommer, R., Spectral sensitivity of *Bacillus subtilis* spores and MS2 Coliphage for validation testing of ultraviolet reactors for water disinfection. *Environmental Science & Technology* **2005**, *39*, (20), 7845-7852.
14. Woo, M. H.; Grippin, A.; Anwar, D.; Smith, T.; Wu, C. Y.; Wander, J. D., Effects of Relative Humidity and Spraying Medium on UV Decontamination of Filters Loaded with Viral Aerosols. *Applied and Environmental Microbiology* **2012**, *78*, (16), 5781-5787.
15. Peccia, J.; Werth, H. M.; Miller, S.; Hernandez, M., Effects of relative humidity on the ultraviolet induced inactivation of airborne bacteria. *Aerosol Science and Technology* **2001**, *35*, (3), 728-740.
16. Fletcher, L. A., The Influence of Relative Humidity on the UV Susceptibility of Airborne Gram Negative Bacteria. *IUVA News* 2004, pp 12-19.

17. Ko, G.; First, M. W.; Burge, H. A., Influence of relative humidity on particle size and UV sensitivity of *Serratia marcescens* and *Mycobacterium bovis* BCG aerosols. *Tubercle and Lung Disease* **2000**, *80*, (4-5), 217-228.
18. Tseng, C. C.; Li, C. S., Inactivation of viruses on surfaces by ultraviolet germicidal irradiation. *Journal of Occupational and Environmental Hygiene* **2007**, *4*, (6), 400-405.
19. Ratnesar-Shumate, S.; Williams, G.; Green, B.; Krause, M.; Holland, B.; Wood, S.; Bohannon, J.; Boydston, J.; Freeburger, D.; Hooper, I.; Beck, K.; Yeager, J.; Altamura, L. A.; Biryukov, J.; Yolitz, J.; Schuit, M.; Wahl, V.; Hevey, M.; Dabisch, P., Simulated Sunlight Rapidly Inactivates SARS-CoV-2 on Surfaces. *Journal of Infectious Diseases* **2020**, *XX*, 1-9.
20. Buonanno, M.; Welch, D.; Shuryak, I.; Brenner, D. J., Far-UVC light (222 nm) efficiently and safely inactivates airborne human coronaviruses. *Scientific Reports* **2020**.